

The Joy of MOIs

UIUC 2024

Eva-Maria Hekkelman

UNSW

October 22 2024

Summary of this talk

- 1 Motivation for studying MOIs
- 2 Pseudodifferential calculus
- 3 MOIs of pseudodifferential operators

This talk is based on joint work with Ed McDonald and Teun van Nuland.

Part 1: Motivation

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

In other words, does $|f(t) - f(s)| \lesssim |t - s|$ imply

$$\|f(A) - f(B)\|_p \lesssim \|A - B\|_p?$$

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

In other words, does $|f(t) - f(s)| \lesssim |t - s|$ imply

$$\|f(A) - f(B)\|_p \lesssim \|A - B\|_p?$$

(Answer: no if $p = 1$ or $p = \infty$, yes if $1 < p < \infty$.)

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

In other words, does $|f(t) - f(s)| \lesssim |t - s|$ imply

$$\|f(A) - f(B)\|_p \lesssim \|A - B\|_p?$$

(Answer: no if $p = 1$ or $p = \infty$, yes if $1 < p < \infty$.)

How do we effectively study the following operators?

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

In other words, does $|f(t) - f(s)| \lesssim |t - s|$ imply

$$\|f(A) - f(B)\|_p \lesssim \|A - B\|_p?$$

(Answer: no if $p = 1$ or $p = \infty$, yes if $1 < p < \infty$.)

How do we effectively study the following operators?

- $f(A) - f(B)$;

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

In other words, does $|f(t) - f(s)| \lesssim |t - s|$ imply

$$\|f(A) - f(B)\|_p \lesssim \|A - B\|_p?$$

(Answer: no if $p = 1$ or $p = \infty$, yes if $1 < p < \infty$.)

How do we effectively study the following operators?

- $f(A) - f(B)$;
- $[f(A), B]$;

Functional analysis

Question (Krein)

Is every Lipschitz function operator Lipschitz?

In other words, does $|f(t) - f(s)| \lesssim |t - s|$ imply

$$\|f(A) - f(B)\|_p \lesssim \|A - B\|_p?$$

(Answer: no if $p = 1$ or $p = \infty$, yes if $1 < p < \infty$.)

How do we effectively study the following operators?

- $f(A) - f(B)$;
- $[f(A), B]$;
- $\frac{d^n}{dt^n} \big|_{t=0} f(A + tB)$.

Exhibit A

Let us now show that if $b \in \cap \text{Dom } L^k R^q$ then $b \in \text{Dom } \delta$. The proof is more subtle than one would expect, because the obvious argument, using

$$|D| = \pi^{-1} \int_0^\infty \frac{D^2}{D^2 + \mu} \mu^{-1/2} d\mu ,$$

requires some care. Indeed, one gets from the above

$$[|D|, b] = \pi^{-1} \int_0^\infty (D^2 + \mu)^{-1} [D^2, b] (D^2 + \mu)^{-1} \mu^{1/2} d\mu .$$

We can replace $[D^2, b]$ by $|D|$, which has the same size, and get

$$\int_0^\infty (D^2 + \mu)^{-2} |D| \mu^{1/2} d\mu = \int_0^\infty (1+t)^{-2} t^{1/2} dt .$$

For this to work, we need to move $[D^2, b]$ in front of the above integral, i.e. use the finiteness of the norm of

$$\int_0^\infty \underbrace{[(D^2 + \mu)^{-1}, [D^2, b]]}_{-(D^2 + \mu)^{-1} [D^2, [D^2, b]] (D^2 + \mu)^{-1}} (D^2 + \mu)^{-1} \mu^{1/2} d\mu .$$

This finiteness follows from:

- 1) $(D^2 + \mu)^{-1} [D^2, [D^2, b]]$ bounded since $b \in \text{Dom } L^2$
- 2) $\int_0^\infty \|(D^2 + \mu)^{-2}\| \mu^{1/2} d\mu \leq C \int_0^1 \mu^{1/2} d\mu + \int_1^\infty \mu^{-3/2} d\mu < \infty$.

Once $[D^2, b]$ is moved in front the above calculation applies.

From [ConnesMoscovici1995]

Exhibit B

Now, onwards with the computation, the first part of which is straightforward:

$$\begin{aligned}
 [\Delta^{-z}, A]B &= \frac{1}{2\pi i} \int \lambda^{-z} [(\lambda - \Delta)^{-1}, A]B \, d\lambda \\
 &= \frac{1}{2\pi i} \int \lambda^{-z} (\lambda - \Delta)^{-1} [\Delta, A] (\lambda - \Delta)^{-1} B \, d\lambda \\
 &= \int \lambda^{-z} (\lambda - \Delta)^{-1} [\Delta, A] B (\lambda - \Delta)^{-1} \, d\lambda \\
 &\quad + \int \lambda^{-z} (\lambda - \Delta)^{-1} [\Delta, A] (\lambda - \Delta)^{-1} [\Delta, B] (\lambda - \Delta)^{-1} \, d\lambda.
 \end{aligned}$$

(In the last step we did two things at once: we commuted B past $(\lambda - \Delta)^{-1}$ and we then used the formula $[S^{-1}, T] = S^{-1}[T, S]S^{-1}$.) The operators $[\Delta, A]$ and $[\Delta, B]$ have orders 1 and 2, respectively.

Before going on, we shall introduce some better notation for our contour integrals.

2.5 Definition. If D_0, \dots, D_p are differential operators on the closed manifold M , then denote by $I_z(D_0, \dots, D_p)$ the integral

$$\frac{1}{2\pi i} \int \lambda^{-z} D_0 (\lambda - \Delta)^{-1} \dots D_p (\lambda - \Delta)^{-1} \, d\lambda$$

(in the integral, copies of $(\lambda - \Delta)^{-1}$ alternate with the operators D_j). The integral converges if $\operatorname{Re}(z) < n$, in the sense we discussed above, and defines an operator on $C^\infty(M)$.

From [Higson2003]

Exhibit C

Theorem 4.2 (Semifinite Odd Local Index Theorem). *Let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ be an odd finitely summable QC^∞ spectral triple with spectral dimension $p \geq 1$. Let $N = [p/2] + 1$ where $[\cdot]$ denotes the integer part, and let $u \in \mathcal{A}$ be unitary. Then*

$$1) \quad sf(\mathcal{D}, u^* \mathcal{D} u) = \frac{1}{\sqrt{2\pi i}} res_{r=(1-p)/2} \left(\sum_{m=1, \text{odd}}^{2N-1} \phi_m^r(Ch_m(u)) \right)$$

where for $a_0, \dots, a_m \in \mathcal{A}$, $l = \{a + iv : v \in \mathbb{R}\}$, $0 < a < 1/2$, $R_s(\lambda) = (\lambda - (1 + s^2 + \mathcal{D}^2))^{-1}$ and $r > 0$ we define $\phi_m^r(a_0, a_1, \dots, a_m)$ to be

$$\frac{-2\sqrt{2\pi i}}{\Gamma((m+1)/2)} \int_0^\infty s^m \tau \left(\frac{1}{2\pi i} \int_l \lambda^{-p/2-r} a_0 R_s(\lambda) [\mathcal{D}, a_1] R_s(\lambda) \cdots [\mathcal{D}, a_m] R_s(\lambda) d\lambda \right) ds.$$

In particular the sum on the right hand side of 1) analytically continues to a deleted neighbourhood of $r = (1-p)/2$ with at worst a simple pole at $r = (1-p)/2$. Moreover, the complex function-valued cochain $(\phi_m^r)_{m=1, \text{odd}}^{2N-1}$ is a (b, B) cocycle for \mathcal{A} modulo functions holomorphic in a half-plane containing $r = (1-p)/2$.

From [CareyPhillipsRennieSukochev2006]

Multiple operator integrals

Let $\phi : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ be such that

$$\phi(\lambda_0, \dots, \lambda_n) = \int_{\Omega} a_0(\lambda_0, \omega) \cdots a_n(\lambda_n, \omega) d\nu(\omega),$$

with finite measure space (Ω, ν) and measurable and bounded $a_j : \mathbb{R} \times \Omega \rightarrow \mathbb{C}$.

Multiple operator integrals

Let $\phi : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ be such that

$$\phi(\lambda_0, \dots, \lambda_n) = \int_{\Omega} a_0(\lambda_0, \omega) \cdots a_n(\lambda_n, \omega) d\nu(\omega),$$

with finite measure space (Ω, ν) and measurable and bounded $a_j : \mathbb{R} \times \Omega \rightarrow \mathbb{C}$.

Let H_0, \dots, H_n be self-adjoint, for $V_1, \dots, V_n \in B(\mathcal{H})$ define the MOI

$$\begin{aligned} T_{\phi}^{H_0, \dots, H_n}(V_1, \dots, V_n)\psi \\ := \int_{\Omega} a_0(H_0, \omega) V_1 a_1(H_1, \omega) \cdots V_n a_n(H_n, \omega) \psi d\nu(\omega), \quad \psi \in \mathcal{H}. \end{aligned}$$

Then,

$$T_{\phi}^{H_0, \dots, H_n} : B(\mathcal{H}) \times \cdots \times B(\mathcal{H}) \rightarrow B(\mathcal{H})$$

and this does not depend on how we represent ϕ (its *symbol*).

Divided differences

Symbols of MOIs encountered in the wild are almost always divided differences, which are defined recursively for $f \in C^n(\mathbb{R})$ as

$$f^{[0]}(\lambda) := f(\lambda);$$

$$f^{[n]}(\lambda_0, \dots, \lambda_n) := \frac{f^{[n-1]}(\lambda_0, \dots, \lambda_{n-1}) - f^{[n-1]}(\lambda_1, \dots, \lambda_n)}{\lambda_0 - \lambda_n},$$

with an appropriate limit if $\lambda_0 = \lambda_n$. In particular,

$$\frac{1}{n!} f^{(n)}(\lambda) = f^{[n]}(\lambda, \dots, \lambda).$$

Example

We can write the JLO cocycle as

$$\begin{aligned} \int_{\Delta_n} \mathrm{Tr}(\eta a_0 e^{-t_0 D^2} [D, a_1] e^{-t_1 D^2} \cdots [D, a_n] e^{-t_n D^2}) dt \\ = \mathrm{Tr}(\eta a_0 T_{f[n]}^{D^2}([D, a_1], \dots, [D, a_n])), \end{aligned}$$

with $f(x) = \exp(-x)$.

Summary

MOIs are a powerful tool in analysis:

Summary

MOIs are a powerful tool in analysis:

- $f(H + V) - f(H) = T_{f^{[1]}}^{H+V,H}(V)$ (analogous to Duhamel's formula);

Summary

MOIs are a powerful tool in analysis:

- $f(H + V) - f(H) = T_{f^{[1]}}^{H+V,H}(V)$ (analogous to Duhamel's formula);
- $[f(H), V] = T_{f^{[1]}}^{H,H}([D, V]);$

Summary

MOIs are a powerful tool in analysis:

- $f(H + V) - f(H) = T_{f^{[1]}}^{H+V, H}(V)$ (analogous to Duhamel's formula);
- $[f(H), V] = T_{f^{[1]}}^{H, H}([D, V]);$
- $\frac{d^n}{dt^n} \big|_{t=0} f(H + tV) = T_{f^{[n]}}^{H, \dots, H}(V, \dots, V),$

each of which has been used to obtain sharp estimates. (Potapov, Sukochev, Skripka, Caspers, Montgomery-Smith, McDonald, Peller, ...)

Summary

MOIs are a powerful tool in analysis:

- $f(H + V) - f(H) = T_{f^{[1]}}^{H+V, H}(V)$ (analogous to Duhamel's formula);
- $[f(H), V] = T_{f^{[1]}}^{H, H}([D, V]);$
- $\frac{d^n}{dt^n} \big|_{t=0} f(H + tV) = T_{f^{[n]}}^{H, \dots, H}(V, \dots, V),$

each of which has been used to obtain sharp estimates. (Potapov, Sukochev, Skripka, Caspers, Montgomery-Smith, McDonald, Peller, ...)

Furthermore, MOIs can systematise operator integral techniques in NCG.

A problem

If you write, like in *The Local Index Formula in Noncommutative Geometry* by Nigel Higson, for a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ and $a \in \mathcal{A}$,

$$\begin{aligned} [D^{-2z}, a] &= \left[\int_{\gamma} \lambda^{-z} (\lambda - D^2)^{-1} d\lambda, a \right] \\ &= \int_{\gamma} \lambda^{-z} [(\lambda - D^2)^{-1}, a] d\lambda \\ &= \int_{\gamma} \lambda^{-z} (\lambda - D^2)^{-1} [D^2, a] (\lambda - D^2)^{-1} d\lambda, \end{aligned}$$

then $[D^2, a] \notin B(\mathcal{H})$, so this is not a standard MOI.

Part 2: Abstract pseudodifferential calculus in the style of Connes–Moscovici, Higson, Guillemin

Pseudodifferential operators

On \mathbb{R}^d , a differential operator $L = \sum_{|\alpha| \leq k} a_\alpha(x) \partial^\alpha$ can be written as

$$L = \mathcal{F}^{-1} \circ M_{p_L} \circ \mathcal{F},$$

where M_{p_L} indicates multiplying with the polynomial $p_L(x, \xi) := \sum_{|\alpha| \leq k} a_\alpha(x) \xi^\alpha$.

Pseudodifferential operators

On \mathbb{R}^d , a differential operator $L = \sum_{|\alpha| \leq k} a_\alpha(x) \partial^\alpha$ can be written as

$$L = \mathcal{F}^{-1} \circ M_{p_L} \circ \mathcal{F},$$

where M_{p_L} indicates multiplying with the polynomial $p_L(x, \xi) := \sum_{|\alpha| \leq k} a_\alpha(x) \xi^\alpha$.

Generally speaking, a pseudodifferential operator of order k on \mathbb{R}^d is an operator of the form $L = \mathcal{F}^{-1} \circ M_{p_L} \circ \mathcal{F}$ where the function p_L is more general, such that

$$L : \mathcal{H}^{s+k,2} \rightarrow \mathcal{H}^{s,2},$$

where

$$\mathcal{H}^{s,2}(\mathbb{R}^n) := \{f \in \mathcal{S}'(\mathbb{R}^n) : (1 - \Delta)^{\frac{s}{2}} f \in L_2(\mathbb{R}^n)\},$$

are Bessel potential Sobolev spaces.

Pseudodifferential operators

On \mathbb{R}^d , a differential operator $L = \sum_{|\alpha| \leq k} a_\alpha(x) \partial^\alpha$ can be written as

$$L = \mathcal{F}^{-1} \circ M_{p_L} \circ \mathcal{F},$$

where M_{p_L} indicates multiplying with the polynomial $p_L(x, \xi) := \sum_{|\alpha| \leq k} a_\alpha(x) \xi^\alpha$.

Generally speaking, a pseudodifferential operator of order k on \mathbb{R}^d is an operator of the form $L = \mathcal{F}^{-1} \circ M_{p_L} \circ \mathcal{F}$ where the function p_L is more general, such that

$$L : \mathcal{H}^{s+k,2} \rightarrow \mathcal{H}^{s,2},$$

where

$$\mathcal{H}^{s,2}(\mathbb{R}^n) := \{f \in \mathcal{S}'(\mathbb{R}^n) : (1 - \Delta)^{\frac{s}{2}} f \in L_2(\mathbb{R}^n)\},$$

are Bessel potential Sobolev spaces.

On a Riemannian manifold M , we can define classes of pseudodifferential operators on $L_2(M)$ that locally look like above.

Sobolev spaces

Given an invertible, positive self-adjoint operator Θ on a separable Hilbert space \mathcal{H} , we can define the ‘Sobolev’ spaces \mathcal{H}^s , $s \in \mathbb{R}$, as the completion of $\text{dom } \Theta^s$ under the norm

$$\|\xi\|_s^2 = \langle \xi, \xi \rangle_s := \langle \Theta^s \xi, \Theta^s \xi \rangle_{\mathcal{H}} = \|\Theta^s \xi\|^2, \quad \xi \in \text{dom } \Theta^s.$$

This forms a Hilbert space. We have continuous embeddings

$$\mathcal{H}^t \subseteq \mathcal{H}^s, \quad s \leq t,$$

because

$$\|\Theta^s \xi\| \leq \|\Theta^{s-t}\|_{\infty} \|\Theta^t \xi\|.$$

We put

$$\mathcal{H}^{\infty} := \bigcap_{s \in \mathbb{R}} \mathcal{H}^s, \quad \mathcal{H}^{-\infty} := \bigcup_{s \in \mathbb{R}} \mathcal{H}^s,$$

and we get for free that \mathcal{H}^{∞} is dense in \mathcal{H} .

Analytic order

Even though Θ itself is an unbounded operator on \mathcal{H} , if we regard it as an operator

$$\Theta : \mathcal{H}^1 \rightarrow \mathcal{H}^0 = \mathcal{H},$$

it is a perfectly good bounded operator:

$$\|\Theta\|_{\mathcal{H}^1 \rightarrow \mathcal{H}^0} = \sup_{\xi: \|\Theta\xi\| \leq 1} \|\Theta\xi\| = 1.$$

Analytic order

Even though Θ itself is an unbounded operator on \mathcal{H} , if we regard it as an operator

$$\Theta : \mathcal{H}^1 \rightarrow \mathcal{H}^0 = \mathcal{H},$$

it is a perfectly good bounded operator:

$$\|\Theta\|_{\mathcal{H}^1 \rightarrow \mathcal{H}^0} = \sup_{\xi: \|\Theta\xi\| \leq 1} \|\Theta\xi\| = 1.$$

We define $\text{op}^r(\Theta)$ for $r \in \mathbb{R}$ as those $T : \mathcal{H}^\infty \rightarrow \mathcal{H}^\infty$ that extend to a bounded operator

$$T : \mathcal{H}^{s+r} \rightarrow \mathcal{H}^s, \quad s \in \mathbb{R}.$$

Analytic order

Even though Θ itself is an unbounded operator on \mathcal{H} , if we regard it as an operator

$$\Theta : \mathcal{H}^1 \rightarrow \mathcal{H}^0 = \mathcal{H},$$

it is a perfectly good bounded operator:

$$\|\Theta\|_{\mathcal{H}^1 \rightarrow \mathcal{H}^0} = \sup_{\xi: \|\Theta\xi\| \leq 1} \|\Theta\xi\| = 1.$$

We define $\text{op}^r(\Theta)$ for $r \in \mathbb{R}$ as those $T : \mathcal{H}^\infty \rightarrow \mathcal{H}^\infty$ that extend to a bounded operator

$$T : \mathcal{H}^{s+r} \rightarrow \mathcal{H}^s, \quad s \in \mathbb{R}.$$

We define $\text{OP}^r(\Theta)$ as those $T \in \text{op}^r(\Theta)$ for which $[\Theta, T] \in \text{op}^r(\Theta)$, $[\Theta, [\Theta, T]] \in \text{op}^r(\Theta)$, $\delta_\Theta^n(T) \in \text{op}^r(\Theta)$.

Examples

- If Δ is the Laplace operator on \mathbb{R}^n , setting $\Theta = (1 - \Delta)^{1/2}$ gives the standard (Bessel potential) Sobolev spaces. The k -th order (pseudo)differential operators are contained in $OP^k(\Theta)$.

Examples

- If Δ is the Laplace operator on \mathbb{R}^n , setting $\Theta = (1 - \Delta)^{1/2}$ gives the standard (Bessel potential) Sobolev spaces. The k -th order (pseudo)differential operators are contained in $OP^k(\Theta)$.
- Taking $\Theta = (1 - \Delta)^{1/2}$ where Δ is the sub-Laplacian on a stratified Lie group gives the Sobolev spaces defined by Folland and Stein.

Examples

- If Δ is the Laplace operator on \mathbb{R}^n , setting $\Theta = (1 - \Delta)^{1/2}$ gives the standard (Bessel potential) Sobolev spaces. The k -th order (pseudo)differential operators are contained in $OP^k(\Theta)$.
- Taking $\Theta = (1 - \Delta)^{1/2}$ where Δ is the sub-Laplacian on a stratified Lie group gives the Sobolev spaces defined by Folland and Stein.
- Can take Θ to be a harmonic oscillator, Kolmogorov operator, or a positive elliptic operator in any kind of pseudodifferential calculus.

Examples

- If Δ is the Laplace operator on \mathbb{R}^n , setting $\Theta = (1 - \Delta)^{1/2}$ gives the standard (Bessel potential) Sobolev spaces. The k -th order (pseudo)differential operators are contained in $OP^k(\Theta)$.
- Taking $\Theta = (1 - \Delta)^{1/2}$ where Δ is the sub-Laplacian on a stratified Lie group gives the Sobolev spaces defined by Folland and Stein.
- Can take Θ to be a harmonic oscillator, Kolmogorov operator, or a positive elliptic operator in any kind of pseudodifferential calculus.
- In NCG, given a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ it makes sense to put $\Theta = (1 + D^2)^{1/2}$. Then for example $D \in OP^1(\Theta)$, and for a *regular* spectral triple $a, [D, a] \in OP^0(\Theta)$ for all $a \in \mathcal{A}$.

Examples

- If Δ is the Laplace operator on \mathbb{R}^n , setting $\Theta = (1 - \Delta)^{1/2}$ gives the standard (Bessel potential) Sobolev spaces. The k -th order (pseudo)differential operators are contained in $OP^k(\Theta)$.
- Taking $\Theta = (1 - \Delta)^{1/2}$ where Δ is the sub-Laplacian on a stratified Lie group gives the Sobolev spaces defined by Folland and Stein.
- Can take Θ to be a harmonic oscillator, Kolmogorov operator, or a positive elliptic operator in any kind of pseudodifferential calculus.
- In NCG, given a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ it makes sense to put $\Theta = (1 + D^2)^{1/2}$. Then for example $D \in OP^1(\Theta)$, and for a *regular* spectral triple $a, [D, a] \in OP^0(\Theta)$ for all $a \in \mathcal{A}$.
- If Θ is bounded, $\mathcal{H}^s \simeq \mathcal{H}$ and $op^r(\Theta) = B(\mathcal{H})$ for all $s, r \in \mathbb{R}$.

Elliptic operators

Our goal is to construct MOIs where all operators are in $\text{op}(\Theta)$. First, we need a functional calculus for such operators. Analogously to usual notions of pseudodifferential operators, a functional calculus can be constructed for *elliptic* operators.

Elliptic operators

Our goal is to construct MOIs where all operators are in $\text{op}(\Theta)$. First, we need a functional calculus for such operators. Analogously to usual notions of pseudodifferential operators, a functional calculus can be constructed for *elliptic* operators.

We define $T \in \text{op}^r(\Theta)$ to be Θ -*elliptic*, if there is a parametrix $P \in \text{op}^{-r}(\Theta)$ such that

$$TP = 1_{\mathcal{H}^\infty} + \text{op}^{-\infty}(\Theta);$$

$$PT = 1_{\mathcal{H}^\infty} + \text{op}^{-\infty}(\Theta).$$

By a Borel Lemma argument, it suffices if

$$TP = 1_{\mathcal{H}^\infty} + \text{op}^{-1}(\Theta);$$

$$PT = 1_{\mathcal{H}^\infty} + \text{op}^{-1}(\Theta).$$

Elliptic operators 2

If $\Theta = (1 + D^2)^{1/2}$, then D is Θ -elliptic, and so is $D + V$ if $V \in \text{op}^r$ with $r < 1$.

Elliptic operators 2

If $\Theta = (1 + D^2)^{1/2}$, then D is Θ -elliptic, and so is $D + V$ if $V \in \text{op}^r$ with $r < 1$.

Let $T \in \text{op}^r(\Theta)$ be Θ -elliptic,

- If $x \in \mathcal{H}^{-\infty}$, then $Tx \in \mathcal{H}^s$ implies that $x \in \mathcal{H}^{s+r}$ (*elliptic regularity*).
- If $T : \mathcal{H}^r \subseteq \mathcal{H}^0 \rightarrow \mathcal{H}^0$ (i.e. $r \geq 0$) is a symmetric operator, then it is self-adjoint. This situation will be referred to as ' T is Θ -elliptic and symmetric'.

Elliptic operators 2

If $\Theta = (1 + D^2)^{1/2}$, then D is Θ -elliptic, and so is $D + V$ if $V \in \text{op}^r$ with $r < 1$.

Let $T \in \text{op}^r(\Theta)$ be Θ -elliptic,

- If $x \in \mathcal{H}^{-\infty}$, then $Tx \in \mathcal{H}^s$ implies that $x \in \mathcal{H}^{s+r}$ (*elliptic regularity*).
- If $T : \mathcal{H}^r \subseteq \mathcal{H}^0 \rightarrow \mathcal{H}^0$ (i.e. $r \geq 0$) is a symmetric operator, then it is self-adjoint. This situation will be referred to as ' T is Θ -elliptic and symmetric'. **Note: this does *not* imply that**

$$T : \mathcal{H}^{r+s} \subseteq \mathcal{H}^s \rightarrow \mathcal{H}^s$$

is self-adjoint for any other $s \in \mathbb{R}$. In fact, these operators need not even be symmetric or normal.

Functional calculus

We write $f \in L^\beta_\infty(\mathbb{R})$ for some $\beta \in \mathbb{R}$ if $f(x)(1+x^2)^{-\beta/2} \in L_\infty(\mathbb{R})$.

Functional calculus

We write $f \in L_\infty^\beta(\mathbb{R})$ for some $\beta \in \mathbb{R}$ if $f(x)(1+x^2)^{-\beta/2} \in L_\infty(\mathbb{R})$.

H.–McDonald–van Nuland (2024)

Let $T \in \text{op}^r(\Theta)$, $r > 0$, be Θ -elliptic and symmetric. If $f \in L_\infty^\beta(\mathbb{R})$, then

$$f(T) \in \text{op}^{r\beta}(\Theta).$$

Functional calculus

We write $f \in L_\infty^\beta(\mathbb{R})$ for some $\beta \in \mathbb{R}$ if $f(x)(1+x^2)^{-\beta/2} \in L_\infty(\mathbb{R})$.

H.–McDonald–van Nuland (2024)

Let $T \in \text{op}^r(\Theta)$, $r > 0$, be Θ -elliptic and symmetric. If $f \in L_\infty^\beta(\mathbb{R})$, then

$$f(T) \in \text{op}^{r\beta}(\Theta).$$

Furthermore, if A is self-adjoint on \mathcal{H} , $A \in \text{op}^t(\Theta)$, $t \in \mathbb{R}$, and A commutes strongly with T , then for $f \in L_\infty^\beta(\mathbb{R})$, $\beta \geq 0$, we have

$$f(A) \in \text{op}^{t\beta}(\Theta).$$

This second part applies for example to $i\frac{d}{dx}$ in $\text{op}(1-\Delta)^{1/2}$ on \mathbb{R}^d .

Part 3: MOIs as pseudodifferential operators

Unbounded MOIs

H.–McDonald–van Nuland (2024)

Let $H_i \in \text{op}^{h_i}(\Theta)$, $h_i > 0$ Θ -elliptic and symmetric for $i = 0, \dots, n$, and $X_i \in \text{op}^{r_i}(\Theta)$ for $i = 1, \dots, n$. Let $\phi : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ such that

$$\phi(\lambda_0, \dots, \lambda_n) = \int_{\Omega} a_0(\lambda_0, \omega) \cdots a_n(\lambda_n, \omega) d\nu(\omega),$$

with finite measure space (Ω, ν) with $a_j(x, \omega)(1 + x^2)^{-\beta_j/2} : \mathbb{R} \times \Omega \rightarrow \mathbb{C}$ measurable and bounded. Then for $\psi \in \mathcal{H}^\infty$,

$$T_\phi^{H_0, \dots, H_n}(X_1, \dots, X_n)\psi := \int_{\Omega} a_0(H_0, \omega) X_1 a_1(H_1, \omega) \cdots X_n a_n(H_n, \omega) \psi d\nu(\omega)$$

is a well-defined vector in \mathcal{H}^∞ independent of the representation of ϕ , and

$$T_\phi^{H_0, \dots, H_n} : \text{op}^{r_1}(\Theta) \times \cdots \times \text{op}^{r_n}(\Theta) \rightarrow \text{op}^{\sum_j r_j + \sum_j \beta_j h_j}(\Theta).$$

Unbounded MOIs: the useful bit

If $f \in C^{n+2}(\mathbb{R})$, and $f^{(k)} \in L_{\infty}^{\beta-k}(\mathbb{R})$ for $k = 0, \dots, n+2$, then for $H \in \text{op}^h(\Theta)$, $h > 0$ Θ -elliptic and symmetric, and $X_i \in \text{op}^{r_i}(\Theta)$,

$$T_{f^{[n]}}^{H, \dots, H}(X_1, \dots, X_n) \in \bigcap_{\varepsilon > 0} \text{op}^{(\beta-n)h + \sum_j r_j + \varepsilon}(\Theta).$$

If $f \in C^{\infty}(\mathbb{R})$ and $f^{(k)} \in L_{\infty}^{\beta-k}(\mathbb{R})$ for all $k \in \mathbb{N}$, we write $f \in S^{\beta}(\mathbb{R})$.

Unbounded MOIs: the useful bit

If $f \in C^{n+2}(\mathbb{R})$, and $f^{(k)} \in L_{\infty}^{\beta-k}(\mathbb{R})$ for $k = 0, \dots, n+2$, then for $H \in \text{op}^h(\Theta)$, $h > 0$ Θ -elliptic and symmetric, and $X_i \in \text{op}^{r_i}(\Theta)$,

$$T_{f^{[n]}}^{H, \dots, H}(X_1, \dots, X_n) \in \bigcap_{\varepsilon > 0} \text{op}^{(\beta-n)h + \sum_j r_j + \varepsilon}(\Theta).$$

If $f \in C^{\infty}(\mathbb{R})$ and $f^{(k)} \in L_{\infty}^{\beta-k}(\mathbb{R})$ for all $k \in \mathbb{N}$, we write $f \in S^{\beta}(\mathbb{R})$.

The integral we saw earlier,

$$\int_{\gamma} \lambda^{-z} (\lambda - D^2)^{-1} [D^2, a] (D^2 - \lambda)^{-1} d\lambda = T_{f^{[1]}}^{D^2, D^2}([D^2, a]),$$

with $f(x) = x^{-z}$.

Two rules

MOIs as we defined them come with two identities:

- ① $f(A) - f(B) = T_{f^{[1]}}^{A,B}(A - B);$
- ② $[f(H), a] = T_{f^{[1]}}^{H,H}([H, a]),$

Two rules

MOIs as we defined them come with two identities:

$$\textcircled{1} \quad f(A) - f(B) = T_{f^{[1]}}^{A,B}(A - B);$$

$$\textcircled{2} \quad [f(H), a] = T_{f^{[1]}}^{H,H}([H, a]),$$

and the higher order analogues (since $T_{f^{[0]}}^H() = f(H)$)

$$\begin{aligned} \textcircled{1} \quad & T_{f^{[n]}}^{H_0, \dots, A, \dots, H_n}(V_1, \dots, V_n) - T_{f^{[n]}}^{H_0, \dots, B, \dots, H_n}(V_1, \dots, V_n) \\ &= T_{f^{[n+1]}}^{H_0, \dots, A, B, \dots, H_n}(V_1, \dots, A - B, \dots, V_n); \end{aligned}$$

$$\begin{aligned} \textcircled{2} \quad & T_{f^{[n]}}^{H_0, \dots, H_n}(V_1, \dots, V_{j-1}, aV_j, \dots, V_n) - T_{f^{[n]}}^{H_0, \dots, H_n}(V_1, \dots, V_{j-1}a, V_j, \dots, V_n) \\ &= T_{f^{[n+1]}}^{H_0, \dots, H_j, H_j, \dots, H_n}(V_1, \dots, V_{j-1}, [H_j, a], V_{j+1}, \dots, V_n). \end{aligned}$$

Taylor expansion

The first rule on its own gives a Taylor expansion:

$$\begin{aligned} f(H + V) &\stackrel{(1)}{=} f(H) + T_{f^{[1]}}^{H+V, H}(V) \\ &\stackrel{(1)}{=} f(H) + T_{f^{[1]}}^{H, H}(V) + T_{f^{[2]}}^{H+V, H, H}(V, V), \end{aligned}$$

and repeat. We get for all $N \in \mathbb{N}$

$$f(H + V) = \sum_{n=0}^N T_{f^{[n]}}^{H, \dots, H}(V, \dots, V) + T_{f^{[N+1]}}^{H+V, H, \dots, H}(V, \dots, V).$$

Note: if H and V commute,

$$T_{f^{[n]}}^{H, \dots, H}(V, \dots, V) = \frac{1}{n!} f^{(n)}(H) V^n.$$

Commutator expansion

In similar manner, by rule (2) we get

$$T_{f^{[n]}}^{H, \dots, H}(V_1, \dots, V_n) = V_1 T_{f^{[n]}}^{H, \dots, H}(1, V_2, \dots, V_n) + T_{f^{[n+1]}}^{H, \dots, H}([H, V], 1, V_2, \dots, V_n),$$

repeating and remembering that $T_{f^{[n]}}^{H, \dots, H}(1, \dots, 1) = \frac{1}{n!} f^{(n)}(H)$, we get

$$\begin{aligned} T_{f^{[n]}}^{H, \dots, H}(V_1, \dots, V_n) &= \sum_{m=0}^N \sum_{m_1 + \dots + m_n = m} \frac{C_{m_1, \dots, m_n}}{(n+m)!} \delta_H^{m_1}(X_1) \cdots \delta_H^{m_n}(X_n) f^{(n+m)}(H) \\ &\quad + S_{H, V}^N. \end{aligned}$$

Commutator expansion

In similar manner, by rule (2) we get

$$T_{f^{[n]}}^{H, \dots, H}(V_1, \dots, V_n) = V_1 T_{f^{[n]}}^{H, \dots, H}(1, V_2, \dots, V_n) + T_{f^{[n+1]}}^{H, \dots, H}([H, V], 1, V_2, \dots, V_n),$$

repeating and remembering that $T_{f^{[n]}}^{H, \dots, H}(1, \dots, 1) = \frac{1}{n!} f^{(n)}(H)$, we get

$$\begin{aligned} T_{f^{[n]}}^{H, \dots, H}(V_1, \dots, V_n) &= \sum_{m=0}^N \sum_{m_1 + \dots + m_n = m} \frac{C_{m_1, \dots, m_n}}{(n+m)!} \delta_H^{m_1}(X_1) \cdots \delta_H^{m_n}(X_n) f^{(n+m)}(H) \\ &\quad + S_{H, V}^N. \end{aligned}$$

The combinatorics to get this expression is exactly the same as how one gets the cocycle of the local index formula, writing $A^{(k)} := \delta_{D^2}^n(A)$,

$$\begin{aligned} &\phi_n(a_0, \dots, a_n) \\ &= \sum_{|k|, q \geq 0} c_{n, k, q} \operatorname{Res}_{z=0} z^q \operatorname{Tr} \left(a_0 [D, a_1]^{(k_1)} \cdots [D, a_n]^{(k_n)} |D|^{-2|k| - 2z - n} \right), \end{aligned}$$

Asymptotic expansions

We say that $T \sim \sum_{k=0}^{\infty} T_k$ for $T, T_k \in \text{op}(\Theta)$ if

$$T - \sum_{k=1}^N T_k \in \text{op}^{m_N}(\Theta), \quad m_N \downarrow -\infty.$$

Asymptotic expansions

We say that $T \sim \sum_{k=0}^{\infty} T_k$ for $T, T_k \in \text{op}(\Theta)$ if

$$T - \sum_{k=1}^N T_k \in \text{op}^{m_N}(\Theta), \quad m_N \downarrow -\infty.$$

If $f \in S^{\beta}(\mathbb{R})$, if $H \in \text{op}^h(\Theta)$, $h > 0$ is Θ -elliptic and symmetric, and if $V \in \text{op}^r(\Theta)$ with $r < h$, then

$$f(H + V) \sim \sum_{n=0}^{\infty} T_{f^{[n]}}^{H, \dots, H}(V, \dots, V).$$

Asymptotic expansions

We say that $T \sim \sum_{k=0}^{\infty} T_k$ for $T, T_k \in \text{op}(\Theta)$ if

$$T - \sum_{k=1}^N T_k \in \text{op}^{m_N}(\Theta), \quad m_N \downarrow -\infty.$$

If $f \in S^{\beta}(\mathbb{R})$, if $H \in \text{op}^h(\Theta)$, $h > 0$ is Θ -elliptic and symmetric, and if $V \in \text{op}^r(\Theta)$ with $r < h$, then

$$f(H + V) \sim \sum_{n=0}^{\infty} T_{f^{[n]}}^{H, \dots, H}(V, \dots, V).$$

With mild assumptions on the commutators $\delta_H^n(V)$,

$$f(H + V) \sim \sum_{n,m=0}^{\infty} \sum_{m_1 + \dots + m_n = m} \frac{C_{m_1, \dots, m_n}}{(n+m)!} \delta_H^{m_1}(V) \dots \delta_H^{m_n}(V) f^{(n+m)}(H).$$

A familiar expansion

Recall that

$$[f(\Theta), X] = T_{f^{[1]}}^{\Theta, \Theta}([\Theta, X]).$$

Therefore, for $X \in \text{OP}^r(\Theta)$, the expansions on the last slide give

$$[f(\Theta), X] \sim \sum_{k=1}^{\infty} \frac{1}{k!} \delta_{\Theta}^k(X) f^{(k)}(\Theta).$$

A familiar expansion

Recall that

$$[f(\Theta), X] = T_{f^{[1]}}^{\Theta, \Theta}([\Theta, X]).$$

Therefore, for $X \in \text{OP}^r(\Theta)$, the expansions on the last slide give

$$[f(\Theta), X] \sim \sum_{k=1}^{\infty} \frac{1}{k!} \delta_{\Theta}^k(X) f^{(k)}(\Theta).$$

In particular,

$$[\Theta^{\alpha}, X] \sim \sum_{k=1}^{\infty} \binom{\alpha}{k} \delta_{\Theta}^k(X) \Theta^{\alpha-k}, \quad \alpha \in \mathbb{C},$$

and

$$[\log(\Theta), X] \sim \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \delta_{\Theta}^k(X) \Theta^{-k},$$

and we have that $[\Theta^{\alpha}, X] \in \text{OP}^{r+\Re(\alpha)-1}(\Theta)$ and $[\log(\Theta), X] \in \text{OP}^{r-1}(\Theta)$.

Asymptotic trace expansions

H.–McDonald–van Nuland (2024)

Let $(\mathcal{A}, \mathcal{H}, D)$ be a regular s -summable spectral triple $((1 + D^2)^{-1/2} \in \mathcal{L}_s)$. Let V self-adjoint and bounded, generated by \mathcal{A} and D . Then as $t \rightarrow 0$,

$$\begin{aligned} & \text{Tr}(f(tD + tV)) \\ &= \sum_{n=0}^N \sum_{m=0}^N \sum_{m_1 + \dots + m_n = m} t^{n+m} \frac{C_{m_1, \dots, m_n}}{(n+m)!} \text{Tr}(\delta_D^{m_1}(V) \dots \delta_D^{m_n}(V) f^{(n+m)}(tD)) \\ &+ O(t^{N+1-s}). \end{aligned}$$

Bonus: Functional calculus for OP

If $A \in \text{op}^r(\Theta)$ is Θ -elliptic and symmetric, then by rule (2) we know that for $f \in L^\beta_\infty(\mathbb{R})$

$$f(A) \in \text{op}^{\beta r}(\Theta),$$

$$[\Theta, f(A)] = T_{f[1]}^{A,A}([\Theta, A]),$$

and similar expressions hold for $\delta_\Theta^n(f(A))$. If $A \in \text{OP}^r(\Theta)$, then we can deduce what the order is of these expressions if $f \in S^\beta(\mathbb{R})$, so that

$$\delta_\Theta^n(f(A)) \in \bigcap_{\varepsilon > 0} \text{op}^{r\beta + \varepsilon}(\Theta).$$

We therefore conclude that $f(A) \in \bigcap_{\varepsilon > 0} \text{OP}^{r\beta + \varepsilon}(\Theta)$.

Thanks

Thank you for your attention!